arXiv:1209.1684v1 [quant-ph] 8 Sep 2012

Quantum isobaric process in coupled system and coupled quantum Brayton cycle

X. L. Huang(黄晓理),[1,*] L. C. Wang(王林成),[2] and X. X. Yi(衣学喜)[2,†]
[1]*School of physics and electronic technology,*
Liaoning Normal University, Dalian, 116029, China
[2]*School of physics and optoelectronic technology,*
Dalian University of Technology, Dalian 116024 China

(Dated: September 11, 2012)

We consider the quantum isobaric process and quantum Brayton cycle for a composite system as working substance. The thermodynamics relations between the total system and the subsystem are studied. We can define two pressures for the working substance of the composite system, one corresponds to the external magnetic field (named F_x), the other conjugates to the coupling strength (named F_y), and as a consequence, we can define two types of quantum Brayton cycle for the composite system. We find that the subsystem also experiences a quantum Brayton cycle in the case of cycle based on F_x, whereas the subsystem is a quantum Otto cycle for the cycle based on F_y. The efficiency for the composite system equals to the one for the subsystem in both cases. However the work done by the total system are usually larger than the sum of the two subsystems. Another interesting result is that for the cycle based on F_y, the subsystem can be a refrigerator while the total system is a heat engine under proper condition.

PACS numbers: 05.70.-a, 07.20.Pe, 03.65.-w, 51.30.+i

I. INTRODUCTION

It is well know that there are four basic thermodynamics processes in classical thermodynamics: adiabatic process, isothermal process, isochoric process and isobaric process. These four processes correspond entropy, temperature, volume and pressure unchanged. The quantum mechanics generalization for these concept is a meaningful work. The discovery of the quantum adiabatic theorem [1, 2] brings the adiabat to quantum region. In recent paper, isothermal and isochoric processes are clarified in quantum case [3, 4]. According to these generalizations and the quantum version of the first law of thermodynamics, the quantum Carnot cycle and quantum Otto cycle can be discussed [5, 6]. Several years later, the last concept, quantum isobaric process, is then defined [7] and all kinds of thermodynamics cycles such as Brayton cycle and Diesel cycle [8, 9] can be studied in quantum domain. The studies of quantum thermodynamics cycle [10–15] are usually focused on whether it can surpass the classical limit on the efficiency and work done during a cycle [3, 4, 16] and how to better the work extraction during a cycle [17]. For example, it is reported that one can extract work from a single heat bath via vanishing quantum coherence [16] and the efficiency of a quantum heat engine can be higher than the classical one due to the effect of squeezed heat bath [18]. These studies can better our understanding of the fundamental concept for thermodynamics, statistical mechanics and quantum mechanics. It brings new insights to basic problems in quantum mechanics and thermodynamics [19, 20].

Recently, coupled quantum system as working substance becomes an interesting topic. Several work have been done in the case of quantum Otto cycle [21–24]. On the one hand, the effects of quantum entanglement on the basic thermodynamics quantities during a cycle are studied and it is shown that the second law of thermodynamics holds in such models [21–23]. On the other hand, the effect of coupling on the cycle is also an attractive problem [24]. An interesting result in a coupled quantum Otto cycle is that in certain case the total coupled system absorbs heat from hot bath and release heat to cold bath whereas the local system absorbs heat at cold bath and releases heat at hot bath while performing a net work [24]. In this paper, motivated by these work, we study the effects of coupling on the isobaric process and quantum Brayton cycle. A coupled spins system is taken as working substance, the relation between the total system and the local system during the cycle is studied and several interesting results are obtained.

This paper is organized as follows. In Sec.II, we first give a brief introduction about the pressure in quantum-mechanical system and quantum Brayton cycle. Based on a single spin in external magnetic field, the main properties of the Brayton cycle are presented. In Sec.III, the detailed analysis for the quantum isobaric process and Brayton cycle with coupled system as work substance is performed. Conclusions are given in Sec.IV.

II. QUANTUM ISOBARIC PROCESS AND QUANTUM BRAYTON CYCLE FOR SPIN-$1/2$ SYSTEM IN AN EXTERNAL MAGNETIC FIELD

In this paper, the concept of pressure in quantum-mechanical system is based on the definition given in Ref.[7] as

$$F = -\sum_n p_n \frac{dE_n}{dL}, \tag{1}$$

where L is the generalized coordinate of the system. This definition is derived from the quantum version of the first law of thermodynamics $dU = \sum_n E_n dp_n + \sum_n p_n dE_n \equiv$

---

[*] Electronic address: huangxiaoli1982@foxmail.com
[†] Electronic address: yixx@dlut.edu.cn



FIG. 1: Force-displacement F-L diagram of a quantum Brayton cycle based on spin systems.

$\text{đ}Q + \text{đ}W = TdS + \sum_n Y_n dy_n$ and an analogy of classical relation between generalized force and generalized coordinate as $Y_n = -\frac{\text{đ}W}{dy_n}$, where T and S refer to temperature and thermodynamics entropy, Y_n is the generalized force and y_n is its conjugate generalized coordinate (dy_n can be seen as the generalized displacement). We should mention that for a one dimensional system, the generalized force is the same as pressure. We first consider a spin-$1/2$ system in an external magnetic field as work substance. The Hamiltonian of the work substance can be written as $H = \frac{1}{2}B\sigma^z$. For a system at thermal equilibrium, choosing the inverse of the magnetic field as the generalized coordinate $L = 1/B$, we can calculate the generalized force for such a system according to Eq.(1) as

$$F = -\frac{\tanh(\frac{\beta}{2L})}{2L^2}, \tag{2}$$

where $\beta = 1/kT$, T is the temperature of the system and k is the Boltzmann constant. Then a quantum isobaric process can be defined as a process with a constant force F, which can be realized by controlling the temperature T (or β) and magnetic field B (or L) according to Eq.(2). We should note that the formula of the pressure for different systems are usually different because we do not have a unique equation of state for quantum-mechanical system. This can also been verified in Sec.III.

A quantum Brayton cycle is a quantum generalization of the classical Brayton cycle [8, 9], which consists two quantum adiabatic processes and two quantum isobaric processes (see Fig.1). Starting from point A, the four stages of a cycle can be depicted as follows: Stage 1: $A \to B$ is an isobaric process, in which the generalized force keeps F_1. The system absorbs heat from the environment and some work is done on the system. In order to ensure the heat is absorbed from the environment, we should have $L_B < L_A$. Stage 2: $B \to C$ is an adiabatic process, where only some work is done by the system. Stage 3: $C \to D$ is almost an inverse process of stage 1 that the generalized force is replaced by F_0. Stage 4: $D \to A$ is another adiabatic process. In the isobaric process the generalized force keeps as constant while in the quantum adiabatic process the entropy of the system is fixed. The entropy of the system is

$$S = -k[\ln(2\cosh\frac{\beta}{2L}) - \frac{\beta}{2L}\tanh\frac{\beta}{2L}]. \tag{3}$$

Hence in the quantum adiabatic process we have TL=const (or β/L=const). Based on this fact, we can get a further equation in quantum adiabatic process from Eq.(2) as

$$FL^2 = \text{const}. \tag{4}$$

On the other hand, the internal energy of the system is

$$U = -\frac{1}{2L}\tanh\frac{\beta}{2L}. \tag{5}$$

Comparing it with Eq.(2) we find $FL = U$. Using these two relations, the basic thermodynamics quantities such as heat transfer, net work done by the system and efficiency during the cycle can be obtained as follows

$$\begin{aligned}
Q_{AB} &= \int_{L_A}^{L_B} \sum_n E_n dp_n \\
&= \sum_n p_n E_n \Big|_{L_A}^{L_B} - \int_{L_A}^{L_B} \sum_n p_n \frac{dE_n}{dL} dL \\
&= U(L_B) - U(L_A) + \int_{L_A}^{L_B} F_1 dL = 2F_1(L_B - L_A), \\
Q_{CD} &= 2F_0(L_C - L_D), \\
W &= Q_{AB} - Q_{CD}, \\
\eta &= 1 - \frac{Q_{CD}}{Q_{AB}} = 1 - \sqrt{\frac{F_0}{F_1}}.
\end{aligned}$$

We can see from this equation that the system absorbs heat in the process $A \to B$ and releases heat in the process $C \to D$. The efficiency is consistent to the classical result $\eta = 1 - (\frac{F_0}{F_1})^{1-1/\gamma}$, where γ is the adiabatic exponent. In this model, we have shown that in the adiabatic process TL=const. Comparing with $TL^{\gamma-1}$=const for the adiabatic process, the adiabatic exponent is $\gamma = 2$. These result can also be confirmed for the coupled system as working substance.

III. QUANTUM ISOBARIC PROCESS AND QUANTUM BRAYTON CYCLE FOR COUPLED SPIN-$1/2$ SYSTEM

The main purpose of this paper is to study of the quantum isobaric process and quantum Brayton cycle for the coupled system as work substance and consider the effect of coupling on the cycle. The Hamiltonian of the working substance under consideration is

$$H = \frac{B}{2}(\sigma_1^z + \sigma_2^z) + J(\sigma_1^+\sigma_2^- + \sigma_1^-\sigma_2^+) \tag{6}$$

where J is the coupling constant between the spins. $J > 0$ and $J < 0$ correspond to the antiferromagnetic and the ferromagnetic case respectively. Here we only consider the antiferromagnetic case, i.e., $J > 0$. The four eigenvalues and corresponding eigenstates can be easily obtained as

$$\begin{aligned}
&E_1=-B, \quad |\psi_1\rangle=|00\rangle,\\
&E_2=-J, \quad |\psi_2\rangle=\frac{1}{\sqrt{2}}(|01\rangle-|10\rangle),\\
&E_3=J, \quad\ \ |\psi_3\rangle=\frac{1}{\sqrt{2}}(|01\rangle+|10\rangle),\\
&E_4=B, \quad\ |\psi_4\rangle=|11\rangle. \qquad (7)
\end{aligned}$$

There are two independent parameters in the Hamiltonian, and hence we need two generalized coordinates to describe the system. Choosing $X=1/B$ and $Y=1/J$ as the generalized coordinates, we can define two generalized forces (or pressures) corresponding to X and Y respectively as

$$F_x=-\frac{\sinh\frac{\beta}{X}}{X^2(\cosh\frac{\beta}{X}+\cosh\frac{\beta}{Y})}, \qquad (8)$$

$$F_y=-\frac{\sinh\frac{\beta}{Y}}{Y^2(\cosh\frac{\beta}{X}+\cosh\frac{\beta}{Y})}. \qquad (9)$$

The quantum isobaric process for this system means either F_x or F_y is fixed. As a result, we will discuss these two different cases respectively. The expression for the entropy is too complex, and we do not give it here, however, according to the invariance of entropy in quantum adiabatic process, we have

$$XT=\text{const},\ \ YT=\text{const},\ \ \frac{X}{Y}=\text{const}. \qquad (10)$$

This means that all the energy-level spacings of the work substance change by the same ratio. If we add another restriction which we have adopted in Sec.II that this ratio equals to the ratio of the temperatures of the two baths just before and after the adiabatic processes, the cycle is reversible. Hence the discussion in this paper focuses on the reversible quantum Brayton cycle. The internal energy of the system can be written as

$$U=-\frac{Y\sinh\frac{\beta}{X}+X\sinh\frac{\beta}{Y}}{XY(\cosh\frac{\beta}{X}+\cosh\frac{\beta}{Y})}=F_xX+F_yY, \qquad (11)$$

which is very useful in the following discussions. We should also note that in the following discussion, we consider the relation between the total system and local system during the cycle. Here, in both cases, the symbol F_{loc} or the concept of local force, is the force corresponding to the local magnetic field B for on of the subsystems.

A. Fixed F_x

We first consider the relation between the composite system and subsystem in quantum isobaric process when the generalized force F_x is fixed. In this process, the generalized coordinate and the temperature are controlled according to Eq.(8) such that F_x is a constant with the generalized coordinate Y as another constant. We will proof that both of the subsystems undergo an isobaric process when the generalized force F_x is fixed and the generalized force for the subsystems equals to $\frac{1}{2}F_x$. As a consequence, the cycle for the subsystem is also quantum Brayton cycle.

Proof: the state of the composite system is a thermal equilibrium state which can be easily obtained as

$$\rho(T)=\sum_n p_n|\psi_n\rangle\langle\psi_n|=\begin{pmatrix} p_4 & 0 & 0 & 0\\ 0 & \frac{1}{2}(p_2+p_3) & \frac{1}{2}(p_3-p_2) & 0\\ 0 & \frac{1}{2}(p_3-p_2) & \frac{1}{2}(p_2+p_3) & 0\\ 0 & 0 & 0 & p_1\end{pmatrix}, (12)$$

where $p_n=\frac{1}{\mathcal{Z}}\exp(-\frac{E_n}{kT})$, and T is the temperature of the composite system. $\mathcal{Z}=\sum_n\exp(-\frac{E_n}{kT})$ is the partition function of the system. For the reduced system, we can get the reduced state by tracing out the other subsystem as

$$\rho_A=\rho_B=\begin{pmatrix} p_4+\frac{1}{2}(p_2+p_3) & 0\\ 0 & p_1+\frac{1}{2}(p_2+p_3)\end{pmatrix}. \qquad (13)$$

This state can be seen as an equilibrium state with a local effective temperature β_{loc} (or T_{loc}) as

$$\beta_{\text{loc}}=X\ln\frac{p_1+\frac{1}{2}(p_2+p_3)}{p_4+\frac{1}{2}(p_2+p_3)}. \qquad (14)$$

According to the definition of generalized force for a single spin in Sec.II, we arrive

$$F_{\text{loc}}=-\frac{1}{2X^2}\tanh(\frac{\beta_{\text{loc}}}{2X}). \qquad (15)$$

By virtue of the formula $\tanh(\frac{1}{2}\ln x)=\frac{x-1}{x+1}$ and the definition for F_x (Eq.(8)) we have

$$F_{\text{loc}}=-\frac{\sinh\frac{\beta}{X}}{2X^2(\cosh\frac{\beta}{X}+\cosh\frac{\beta}{Y})}=\frac{1}{2}F_x. \qquad (16)$$

This result shows that if the composite system keeps the generalized force F_x unchanged, the generalized force for subsystem are also invariable and it equals to $\frac{1}{2}F_x$.

In the following, we consider the quantum Brayton cycle based on the coupled system when F_x is fixed. Similar to the case for a single spin, we decreases X in stage 1 and adjust the temperature T so that F_{x1} keeps unchanged. Moreover, the parameters Y_1, which is the inverse of the coupling constant, is also invariable in this stage. In the adiabatic process, Eq.(10) should be satisfied so that the cycle is reversible. After this stage, the parameter becomes Y_0 and in stage 3 it keeps as a constant. The force in stage 3 is F_{x0}. The heat absorbed by the system during the stage 1 is

$$\begin{aligned}
Q_{AB}&=\int_{X_A}^{X_B}\sum_n E_n dp_n=U(X_B)-U(X_A)+\int_{X_A}^{X_B}F_{x1}dX\\
&=2F_{x1}(X_B-X_A)+(F_{yB}-F_{yA})Y_1. \qquad (17)
\end{aligned}$$

We can see from the above expression that Q_{AB} depends not only on the force F_x but also on the force F_y and the coupling constant. This means that the interaction between the



FIG. 2: Efficiency and net work done by the coupled system as a function of J/B at point A when (a) initial pressure F_{x1} (b) initial temperature kT_a is fixed. Other parameters in the figure are $F_{x0} = \frac{1}{4}F_{x1}$, $\frac{X_A}{X_B} = \frac{X_D}{X_C} = 3$.

two spins has a strong effect on the cycle. Similarly, we can calculate the heat released to environment in stage 3 as

$$Q_{CD} = 2F_{x0}(X_C - X_D) + (F_{yC} - F_{yD})Y_0. \tag{18}$$

Hence the efficiency of the cycle can be expressed as

$$\eta = 1 - \frac{Q_{CD}}{Q_{AB}} = 1 - \sqrt{\frac{F_{x0}}{F_{x1}}}. \tag{19}$$

The detailed derivation about the above equation is given in appendix A. This result is the same to the single spin case. After the same procedure as given in Sec.II, we can get the efficiency for the subsystem as

$$\eta_{\text{loc}} = 1 - \sqrt{\frac{F_{\text{loc}0}}{F_{\text{loc}1}}} = 1 - \sqrt{\frac{F_{x0}}{F_{x1}}}, \tag{20}$$

This equation shows that the efficiency for the composite system as working substance is the same to the one for its subsystem. The net work done by the composite system is

$$\begin{aligned} W = {} & [2F_{x1}(X_B - X_A) - 2F_{x0}(X_C - X_D)] \\ & + [(F_{yB} - F_{yA})Y_1 - (F_{yC} - F_{yD})Y_0]. \end{aligned} \tag{21}$$

The first $[\cdot]$ on the right hand side can be seen as the sum of the work done by the two subsystem while the second $[\cdot]$ comes from the effects of interaction between the two subsystem. Due to the positivity of this term, we have $W > 2W_{\text{loc}}$, i.e., the total work performed is larger than the sum of work obtained from the two spins locally. Numerical examples for this result is shown in Fig.2. From the figure we confirm that the coupling can increase the net work done by the system during a cycle although the efficiency does not been improved. When $J = 0$, $W/W_{\text{loc}} = 2$ in all cases, as expected. Another point to be explained is that in Fig.2, some lines are interruptive. This is because that when the absolute value of F_{x1} is larger, the possible region for the coupling constant is smaller.

B. Fixed F_y



FIG. 3: The local heat exchange Q_1 and Q_2 and the local work W_{loc} (in units of B) as a function of J/B. Here the initial temperature $kT_a = 0.5B$. Other parameters are $F_{y0} = \frac{1}{4}F_{y1}$, $\frac{Y_A}{Y_B} = \frac{Y_D}{Y_C} = 3$.

Similar to the discussion given above, an isobaric process that F_y is fixed means one should carefully control the coordinate Y and temperature T such that F_y is a constant with X as another constant. It can be easily verified that during this process, the pressure for the subsystem is not fixed, i.e., the process for the subsystem is not an isobaric process. However, when we control the coordinate Y for the composite system, the local Hamiltonian for the subsystem does not vary and local energy levels keep as constants. As a result, the subsystems undergo quantum isochoric process when F_y is fixed, and a quantum Brayton cycle based on F_y for the composite system corresponds to a quantum Otto cycle for the subsystem.

After some calculation which is similar to Sec.III A, we obtain the efficiency of the cycle for the composite system as

$$\eta_y = 1 - \sqrt{\frac{F_{y0}}{F_{y1}}}. \tag{22}$$

The efficiency for the subsystem can be obtained according to the efficiency of quantum Otto cycle as

$$\eta_{\text{loc}} = 1 - \frac{X_1}{X_0} = 1 - \sqrt{\frac{F_{y0}}{F_{y1}}}. \tag{23}$$

This result tells us that the efficiency for the composite system during a quantum Brayton cycle based on F_y equals to the one for the subsystem during a quantum Otto cycle. The precondition for this conclusion is that the subsystem is a Otto heat engine. We should note that when initial temperature is small enough and the coupling constant is large enough, the work done by the subsystem can be negative. In this case, the subsystem is a refrigerator while the total system is a heat engine. Numerical examples are given in Fig.3. In this figure we calculate the heat exchange Q_1 in stage 1 and Q_2 in stage 3 for the subsystem. Here $Q_i > 0$ and $Q_i < 0$ $(i = 1, 2)$ correspond to absorption and release of heat, W_{loc} is the local work. From the figure we see that when initial J/B at point A is larger than a certain value, $Q_1 < 0$, $Q_2 > 0$ and $W_{\text{loc}} < 0$. This indicates that the subsystem is a refrigerator, it releases heat when the composite system absorbs heat in stage 1 and vice versa in stage 3. The main reason for this phenomenon comes from the strange property for the eigenstate $|\psi_3\rangle$ of the system. In the strong coupling region, the ground state of the system is $|\psi_3\rangle$. However, the reduced state for this state is $\rho_{A(B)} = \frac{1}{2}I_2$, which can be seen as an equilibrium state at local effective temperature $T_{\text{loc}} = \infty$. Here I_2 is a 2×2 unit matrix. Hence in stage 1, the globe temperature increases while the local temperature decreases when the globe temperature is low enough and the coupling constant is large enough. Hence the subsystem releases heat in stage 1 and inversely in stage 3. As a result the local cycle is a refrigeration cycle.

IV. CONCLUSION

In summary, we have studied the quantum isobaric process and quantum Brayton cycle in which coupled spins system is taken as working substance. The concept of force or pressure in quantum system is derived from the quantum version of the first law of thermodynamics and an analogy of classical relation between generalized force and generalized coordinate, which is first proposed in Ref.[7]. A quantum Brayton cycle consists of two quantum adiabatic processes and two quantum isobaric processes. There are two generalized coordinates in the system, i.e., the local external magnetic field and the coupling strength, therefore we can define two pressures respectively and construct two types of quantum Brayton cycle. We find that in quantum Brayton cycle based on the pressure corresponding to the external field, the subsystem undergo a quantum Brayton cycle with pressure half of the total system, while in the cycle based on the force conjugating to the coupling strength, the subsystem experiences a quantum Otto cycle. The efficiency for the coupled system in the two cycles are equal to the one for the subsystem, which is formally consistent to the classical result. However the net work done by the total system are usually more than the sum of subsystems. Moreover, when the initial temperature and initial coupling strength are chosen properly, an interesting phenomenon can be observed in the Brayton cycle based on the force corresponding to coupling constant that the total system performed as heat engine while the local spins serve as a refrigerator.

This work is supported by NSF of China under Grant Nos. 11105064, 10905007, and 11175032.

Appendix A: The derivation of η for the coupled system when F_x is fixed.

First, from expression of F_x in Eq.(8) and the condition for adiabatic process in Eq.(10), we have

$$F_x X^2 = \text{const} \tag{A1}$$

As a result, one can construct the relation during the two adiabatic processes as

$$\frac{F_{x0}}{F_{x1}} = \frac{X_B^2}{X_C^2} = \frac{X_A^2}{X_D^2}. \tag{A2}$$

In the similar manner, for the force F_y we obtain

$$F_y Y^2 = \text{const} \tag{A3}$$

and

$$\frac{Y_1^2}{Y_0^2} = \frac{F_{yC}}{F_{yB}} = \frac{F_{yD}}{F_{yA}}. \tag{A4}$$

Moreover, based on $\frac{X}{Y} = \text{const}$ in the adiabatic process, we have

$$\frac{Y_1}{Y_0} = \frac{X_A}{X_D} = \frac{X_B}{X_C} \tag{A5}$$

Combine Eqs.(A2), (A4) and (A5), we obtain a complete ratio relation for the whole cycle as

$$\frac{Y_1^2}{Y_0^2} = \frac{X_A^2}{X_D^2} = \frac{X_B^2}{X_C^2} = \frac{F_{x0}}{F_{x1}} = \frac{F_{yC}}{F_{yB}} = \frac{F_{yD}}{F_{yA}}. \tag{A6}$$

Using this equation we can obtain Eq.(19). In a similar way Eq.(22) can be obtained.

---

[1] M. Born and V. Fork. Z. Phys. 51, 165 (1928).
[2] A. Messiah, *Quantum Mechanics* (Dover, New York, 1999).
[3] T. D. Kieu, Phys. Rev. Lett. **93**, 140403 (2004).
[4] T. D. Kieu, Eur. Phys. J. D **39**, 115 (2006); quan-ph/0311157.
[5] H. T. Quan. P. Zhang, and C. P. Sun, Phys. Rev. E **72**, 056110 (2005).
[6] H. T. Quan, Y. X. Liu, C. P. Sun, and F. Nori, Phys. Rev. E **76**, 031105 (2007).
[7] H. T. Quan, Phys. Rev. E **79**, 041129 (2009).
[8] H. B. Callen, *Thermodynamics and an Introduction to Themostatistics*, 2nd ed. (Wiley, New York, 1985); C. Kittle and H. Kroemer, *Thermal Physics*, 2nd ed. (Freeman, San Francisco, 1980).
[9] P. Perrot, *A to Z of Thermodynamics* (Oxford University Press, Oxford, 1998).
[10] O. Fialko and D. W. Hallwood, Phys. Rev. Lett. **108**, 085303 (2012).
[11] J. Z. He, J. C. Chen, and B. Hua, Phys. Rev. E **65**, 036145 (2002).
[12] T. Feldmann and R. Kosloff, Phys. Rev. E **61**, 4774 (2000); Phys. Rev. E **68**, 016101 (2003); Phys. Rev. E **70**, 046110 (2004).
[13] F. Wu, L. G. Chen, S. Wu, F. R. Sun, C. Wu, J. Chem. Phys. **124**, 214702 (2006).
[14] B. H. Lin and J. C. Chen, Phys. Rev. E **67**, 046105 (2003).
[15] S. Li, H. Wang, Y. D. Sun, and X. X. Yi, J. Phys. A **40**, 8655 (2007).
[16] M. O. Scully, M. S. Zubairy, G. S. Agarwal, and H. Walther, Science **299**, 862 (2003).
[17] J. H. Wang, H. Z. He, and X. He, Phys. Rev. E **84**, 041127

(2011); J. H. Wang, H. Z. He, and Z. Q. Wu, Phys. Rev. E **85**, 031145 (2012); J. H. Wang, Z. Q. Wu, and J. Z. He, Phys. Rev. E **85**, 041148 (2012); J. H. Wang and J. Z. He, J. Appl. Phys. **111**, 043505 (2012).

[18] X. L. Huang, Tao Wang, and X. X. Yi, submitted.

[19] M. O. Scully, Phys. Rev. Lett. **87**, 220601 (2001); Phys. Rev. Lett. **88**, 050602 (2002).

[20] H. T. Quan, Y. D. Wang, Y. X. Liu, C. P. Sun, and F. Nori, Phys. Rev. Lett. **97**, 180402 (2006); H. T. Quan. P. Zhang, and C. P. Sun, Phys. Rev. E **73**, 036122 (2006).

[21] T. Zhang, W. T. Liu, P. X. Chen, and C. Z. Li, Phys. Rev. A **75**, 062102 (2007).

[22] H. Wang, S. Q. Liu, and J. Z. He, Phys. Rev. E **79**, 041113 (2009).

[23] G. F. Zhang, Eur. Phys. J. D **49**, 123 (2008).

[24] G. Thomas and R. S. Johal, Phys. Rev. E **83**, 031135 (2011).